

# Please Keep Your Child Home For . .

## EXCLUSION GUIDELINES FOR CHILD CARE AND EARLY LEARNING FACILITIES



### Vomiting

Exclude a child who vomits two or more times in 24 hours should be excluded until symptoms resolve.



### Diarrhea

Exclude for 2 loose/watery stools more than normal for the child in 24 hours or blood or mucus in the stool.



### Rash, Sores, Lice, Scabies, Ringworm

Exclude for body rash not related to allergy, heat, or diapering; oozing open sores; mouth sores with drooling; or untreated head lice, ringworm or scabies.



### Fever

(with another symptom)

Exclude for fever by any method of 101°F (or 100.4°F for infants less than 2 months) **AND** sore throat, earache, headache, rash, vomiting diarrhea, or just not feeling well.



### Not Feeling Well

Exclude if child is unusually tired, has a loss of appetite, is confused, irritable, pale, or not able to fully participate comfortably.

### When your child is sick:

- 1) Have plans for a back up caregiver for your child
- 2) Call and tell your school or child care what is wrong with your child, even if your child stays home

For a complete list of exclusion guidelines, see WAC 110-300-0205.



CHILD CARE HEALTH OUTREACH PROGRAM  
3020 RUCKER AVENUE, STE. 104  
EVERETT, WA 98201  
(425) 252-5415 [CHILD CARE HEALTH@SNOHD.ORG](mailto:CHILD CARE HEALTH@SNOHD.ORG)