




# INFLUENZA AND PNEUMONIA DEATH

Snohomish County, Washington State & US, 2006-2015

Influenza and pneumonia were the ninth-leading cause of death in Snohomish County in 2015.



Year	SNOHOMISH				WASHINGTON		
	Cases	Population	Age-Adjusted Rate per 100,000	95% C. I.	Cases	Population	Age-Adjusted Rate per 100,000
2006	74	676126	12.9	(10.1, 16.3)	810	6420258	12.8
2007	67	689314	11.3	(8.7, 14.4)	754	6525086	11.5
2008	86	699330	14.3	(11.4, 17.8)	786	6608244	11.8
2009	66	705894	10.2	(7.8, 13.1)	731	6672159	10.7
2010	53	713335	8.0	(5.9, 10.5)	582	6724540	8.3
2011	60	717000	9.0	(6.8, 11.7)	730	6767900	10.3
2012	69	722900	10.7	(8.3, 13.6)	725	6817770	10.0
2013	63	730500	9.8	(7.5, 12.6)	760	6882400	10.1
2014	64	741000	9.0	(6.9, 11.6)	727	6968170	9.5
2015	64	757600	8.4	(6.4, 10.8)	851	7061410	10.7

95% Confidence Interval: A confidence interval is a range of values that has a specified probability of containing the rate. Narrow confidence intervals indicate that rate is precisely estimated, while wider confidence intervals indicate that the estimated rate is less precisely estimated.