




## CHRONIC LOWER RESPIRATORY DISEASE DEATH

Snohomish County, Washington State & US, 2006-2015

Chronic lower respiratory disease was the fifth-leading cause of death in Snohomish County in 2015.



SNOHOMISH					WASHINGTON		
Year	Cases	Population	Age-Adjusted Rate per 100,000		Cases	Population	Age-Adjusted Rate per 100,000
			95% C. I.	Rate per 100,000			
2006	240	676126	(39.1, 50.8)	44.7	2647	6420258	43.4
2007	236	689314	(38.0, 49.4)	43.4	2675	6525086	42.7
2008	226	699330	(34.2, 44.8)	39.2	2933	6608244	45.5
2009	273	705894	(41.0, 52.4)	46.4	2934	6672159	44.5
2010	268	713335	(39.4, 50.5)	44.7	2726	6724540	40.3
2011	283	717000	(39.8, 50.7)	45.0	3081	6767900	44.7
2012	293	722900	(40.4, 51.3)	45.6	2952	6817770	41.5
2013	279	730500	(36.5, 46.6)	41.3	2908	6882400	39.5
2014	280	741000	(35.7, 45.5)	40.3	2907	6968170	38.1
2015	298	757600	(35.8, 45.3)	40.3	3151	7061410	39.9

95% Confidence Interval: A confidence interval is a range of values that has a specified probability of containing the rate. Narrow confidence intervals indicate that rate is precisely estimated, while wider confidence intervals indicate that the estimated rate is less precisely estimated.